

A Dynamical Approach to Gauge Fixing

F. Loran*

Department of Physics, Isfahan University of Technology (IUT)
Isfahan, Iran,
Institute for Studies in Theoretical Physics and Mathematics (IPM)
P. O. Box: 19395-5531, Tehran, Iran.

Abstract

We study gauge fixing in the generalized Gupta-Bleuler quantization. In this method physical states are defined to be simultaneous null eigenstates of a set of quantum invariants. We apply the method to a solvable model proposed by Friedberg, Lee, Pang and Ren and show that no Gribov-type copies appears by construction.

Gauge fixing is significant in quantization of gauge theories. Gribov ambiguities and specially Gribov-type copies [1] have been a challenge for standard methods of quantization. There are some methods to avoid Gribov ambiguities. For example Gribov proposed to restrict field configurations only to those with positive Faddeev-Popov determinant. Recently Klauder [2] has formulated a new method of quantization based on projection operators into gauge invariant states that avoids any gauge fixing procedure and consequently Gribov ambiguities. In this article we propose an alternative approach to this problem. Using the concept of quantum invariants [3], we show that one can avoid Gribov ambiguities by quantizing the whole phase space and then determining the physical (reduced) Hilbert space instead of constructing the Hilbert space based on the reduced phase space [4].

Considering a gauge model given by a Hamiltonian $H(q, p)$ and a set of first class constraints, there exist a generator of gauge transformation that satisfies the following conditions [5, 6],

$$\{G, \phi_\mu^0\}|_{M_0} = 0,$$

*e-mail: loran@cc.iut.ac.ir

$$\left(\{G, H\} + \frac{\partial}{\partial t} G \right)_{M_0} = 0,$$

where $\{ , \}$ stands for the Poisson brackets and M_0 is the submanifold locally defined by primary (first class) constraints ϕ_μ^0 . Gauge transformations $\delta q = \{q, G\}$ and $\delta p = \{p, G\}$, determine gauge orbits i.e. gauge orbits are the integral curves of the generator of gauge transformation. All points on a gauge orbit correspond to physically equivalent configurations of the system. To eliminate gauge freedom (gauge fixing) one may introduce gauge fixing conditions $\chi_i = 0$ that intersect the gauge orbits. A necessary condition for each χ_i to intersect the gauge orbits is that it should not belong to the set of integral curves of G i.e. $\{\chi, G\} \neq 0$ [7, 8, 9]. As is shown formally in Fig.1, a gauge fixing condition χ that satisfies the condition $\{\chi, G\} \neq 0$ may never intersect some of the gauge orbits or there may exist gauge equivalent copies. In these cases, χ 's do not fix (completely) the gauge and Gribov ambiguities emerge [1]. Obviously, Gribov ambiguities would not emerge if one was able to select a certain point on the gauge orbit by determining its coordinates. This can be achieved by using the concept of invariants. It is well known [10] that the classical dynamics of a system possessing gauge degrees of freedom is given by a total Hamiltonian

$$H_T = H + v_\mu \phi_\mu^0, \quad (1)$$

where v_μ 's are Lagrange multipliers. Therefore gauge degrees of freedom are those coordinates q 's which their acceleration $\ddot{q}$ depend on undetermined Lagrange multipliers $v(t)$. It can be easily verified that corresponding to each gauge degree of freedom q there exist a dynamical invariant I_q defined by the relation

$$\frac{\partial}{\partial t} I_q + \{I_q, H_T\} = 0. \quad (2)$$

Consequently the conditions $I_q = 0$ eliminate gauge freedom by determining a point on the gauge orbit. To do quantization one introduces the commutator algebra e.g. $\{q, p\} \rightarrow \frac{1}{i} [\hat{q}, \hat{p}]$ and defines physical states (reduced Hilbert space) as the simultaneous null eigenstates of quantum invariants $\hat{I}_i$ ¹. This method is introduced in our previous work and is called the generalized Gupta-Bleuler quantization [4]. In general the form of the Hamiltonian defined in the generalized Gupta-Bleuler quantization is not the same as the operator form of the total Hamiltonian H_T . Fortunately in the case of important models like Yang-Mills theory and the solvable model proposed by Friedberg, Lee, Pang and Ren [12] these two forms are the same.

¹It is important to note that the eigenvalues of a quantum invariant are independent of time and the particular solutions of the Schrödinger equation are different from the eigenstates of $\hat{I}_i$ only by time dependent phase factors [11]

In the following we study the solvable model proposed by Friedberg, Lee, Pang and Ren. This model can exhibit Gribov copies and is given by a $U(1)$ gauge invariant Lagrangian

$$L = \frac{1}{2} \left[(\dot{x} + gy\xi)^2 + (\dot{y} - gx\xi)^2 + (\dot{z} - \xi)^2 \right] - V \left(\sqrt{x^2 + y^2} \right), \quad (3)$$

or equivalently in terms of polar coordinates [13]

$$L = \frac{1}{2} \dot{r}^2 + \frac{1}{2} r^2 (\dot{\theta} - g\xi)^2 + \frac{1}{2} (\dot{z} - \xi)^2 - V(r). \quad (4)$$

The corresponding total Hamiltonian is

$$H_T = \frac{1}{2} \left(p_r^2 + \frac{L_z^2}{r^2} + p_z^2 \right) + \xi (p_z + gL_z) + V(r) + \ddot{\epsilon}(t) p_\xi, \quad (5)$$

where p_r , L_z , p_z and p_ξ are momenta respectively conjugate to coordinates r , θ , z , ξ and $\ddot{\epsilon}(t)$ is the Lagrange multiplier. Following the generalized Gupta-Bleuler quantization the expectation value of the coordinate operators $q_c = \langle \hat{q} \rangle_{\text{phys}}$ should satisfy the Euler-Lagrange equations of motion obtained from the Lagrangian given in Eq.(4). This can be achieved by introducing the Hamiltonian $\hat{H}^{(0)} = H_T(\hat{q}, \hat{p})$, assuming the commutator algebra $\{ , \} \rightarrow \frac{1}{i} [,]$ and the Schrödinger equation

$$i \frac{\partial}{\partial t} |\text{phys}\rangle = \hat{H}^{(0)} |\text{phys}\rangle. \quad (6)$$

In addition one should impose the condition

$$(\hat{p}_z + g\hat{L}_z) |\text{phys}\rangle = 0 \quad (7)$$

to guarantee the validity of the Lagrangian constraint

$$L_\xi \rightarrow ((\dot{z}_c - \xi_c) + gr_c^2(\dot{\theta}_c - g\xi_c)) = 0.$$

Now for convenience we identify $|\text{phys}\rangle$ as

$$|\text{phys}\rangle = |\psi_\xi\rangle |\psi_z\rangle |\psi_\theta\rangle |\psi_r\rangle. \quad (8)$$

It should be noted that our final result does not depend on this particular form of $|\text{phys}\rangle$. One can easily verify that the operator

$$\hat{I}_\xi = \hat{\xi} - \dot{\epsilon}(t), \quad (9)$$

is an invariant. This means that $|\psi_\xi\rangle$ is the null eigenstate of $\hat{I}_\xi$:

$$\hat{I}_\xi |\psi_\xi\rangle = 0,$$

or equivalently

$$\langle p_\xi | \psi_\xi \rangle = \exp(-ip_\xi \epsilon(t)), \quad (10)$$

where $\hat{p}_\xi |p_\xi\rangle = p_\xi |p_\xi\rangle$. Consequently, the Schrödinger equation Eq.(6) can be written as

$$i \frac{\partial}{\partial t} |\psi_z\rangle |\psi_\theta\rangle |\psi_r\rangle = \hat{H}^{(1)} |\psi_z\rangle |\psi_\theta\rangle |\psi_r\rangle, \quad (11)$$

where $\hat{H}^{(1)}$ is defined as follows,

$$\hat{H}^{(1)} = \frac{1}{2} \left(\hat{p}_r^2 + \frac{\hat{L}_z^2}{\hat{r}^2} + \hat{p}_z^2 \right) + \dot{\epsilon}(t) (\hat{p}_z + g\hat{L}_z) + V(\hat{r}).$$

This Hamiltonian leads to another quantum invariant $\hat{I}_z = \hat{z} - \epsilon(t) - \hat{p}_z t$. This determines $|\psi_z\rangle$ to be

$$\langle p_z | \psi_z \rangle = \exp \left[-i \left(p_z \epsilon(t) + \frac{1}{2} p_z^2 t \right) \right], \quad (12)$$

and the Schrödinger equation takes a new form

$$i \frac{\partial}{\partial t} |\psi_\theta\rangle |\psi_r\rangle = \hat{H}^{(2)} |\psi_\theta\rangle |\psi_r\rangle, \quad (13)$$

where $\hat{H}^{(2)}$ is

$$\hat{H}^{(2)} = \frac{1}{2} \left(\hat{p}_r^2 + \frac{\hat{L}_z^2}{\hat{r}^2} \right) + g\dot{\epsilon}(t)\hat{L}_z + V(\hat{r}).$$

Since $\hat{L}_z$ commutes with $\hat{H}^{(2)}$ one can identify $|\psi_\theta\rangle$ as eigenstates of $\hat{L}_z$;

$$\langle L_z | \psi_\theta \rangle = \exp(-igL_z \epsilon(t)), \quad (14)$$

where L_z is an integer due to periodicity condition on $|\psi_\theta\rangle$. Inserting Eq.(14) in Eq.(13) one obtains the true Schrödinger equation for $|\psi_r\rangle$:

$$i \frac{\partial}{\partial t} |\psi_r^n\rangle = \left[\frac{1}{2} \left(\hat{p}_r^2 + \frac{n^2}{\hat{r}^2} \right) + V(\hat{r}) \right] |\psi_r^n\rangle, \quad (15)$$

where $n = L_z$ is an integer. Therefore the physical states are

$$\langle p_\xi, p_z, L_z, r | \text{phys} \rangle = \exp \left[-i \left(p_\xi \epsilon(t) + p_z \epsilon(t) + \frac{1}{2} p_z^2 t + gL_z \epsilon(t) \right) \right] \psi^n(r). \quad (16)$$

It should be noted that the condition given in Eq.(7) simply relates the eigenvalues of $\hat{p}_z$ and $\hat{L}_z$ to each other i.e. $p_z + gL_z = 0$. Using Eq.(16), one verifies that

$$\begin{aligned} \xi_c &= \dot{\epsilon}(t), \\ z_c &= \epsilon(t) + p_z t \end{aligned}$$

and

$$\theta_c = g\epsilon(t).$$

This means that gauge freedom is properly fixed with no ambiguity. This result can be compared with that of refs.[12, 13]. In one sense our method is similar to the method defined in ref.[2]; in both methods one introduces projection operators into (physical) gauge invariant states. In principle this method can be applied to Yang-Mills theory but first, one should find the clear form of invariants which of course is not a straightforward task.

Acknowledgement

The author would like to thank M. Haghghat and K. Samani for useful discussions and comments.




Fig.1

References

- [1] V. N. Gribov, Nucl. Phys. **B139**, (1978) 1.
- [2] J. R. Klauder, Ann. Phys. **254**, (1997) 419; Nucl. Phys. **B547**, (1999) 397.
- [3] H. R. Lewis and W. B. Riesenfeld, J. Math. Phys. **10**, (1969) 1458.
- [4] F. Loran, hep-th/0203117.
- [5] C. Batlle, J. Gomis, X. Gracia and J. M. Pons, J. Math. Phys. **30** (6), (1986) 1345.
- [6] J. M. Pons and J. A. Garcia, Int. J. Mod. Phys. A **15** (2000) 4681.
- [7] M. Henneaux and C. Teitelboim *"Quantization of Gauge System"* Princeton University Press, Princeton, New Jersey, 1992.
- [8] J. Govaerts, *Hamiltonian Quantization and Constrains Dynamics*, Leuven Notes in Mathematical and Theoretical Physics, Vol. 4, 1991.
- [9] A. Shirzad and F. Loran, hep-th/9912289.
- [10] P. A. M. Dirac, Can. J. Math. **2**, (1950) 129 ; Proc. R. Soc. London Ser. A **246**, (1958) 326; *"Lectures on Quantum Mechanics"* New York: Yeshiva University Press, 1964,
- [11] A. Mostafazadeh, *Dynamical Invariants, Adiabatic Approximation and the Geometric Phase*, Nova Science Publishers, New York, 2001.
- [12] R. Friedberg, T. D Lee, Y. Pang and H. C. Ren, Ann. Phys. **246**, (1996) 381.
- [13] V. M. Villanueva, J. Govaerts, J-L. Lucio-Martinez, J.Phys. A**33**, (2000) 4183.